



## EJVES Extra Abstracts<sup>☆</sup>

### Delayed Presentations of False Aneurysms of the Sub-clavian Artery

N. Haldipur<sup>a,\*</sup>, K. Nagpal<sup>b</sup>, P. Sommaya<sup>a</sup>, P.K. Kumar<sup>c</sup>, J.D. Beard<sup>a</sup>

<sup>a</sup> Vascular Institute, Northern General Hospital, Herries Road, Sheffield, UK

<sup>b</sup> Doncaster Royal Infirmary, Doncaster, UK

<sup>c</sup> Chesterfield Royal Hospital, Chesterfield, UK

Delayed presentation of a false aneurysm following blunt trauma is rare with only five cases discussed in literature.

We describe two cases of late false aneurysms of the Subclavian artery following blunt injury.

doi:10.1016/j.ejvs.2009.01.021

DOI of original article:10.1016/j.ejvsextra.2009.01.002

Available online 26 March 2009

### Percutaneous Angioplasty of the Radial Artery and Its Deep Palmar Branch for Critical Hand Ischemia – A Case Report

A. Cremonesi<sup>\*</sup>, E. Carvalho de Campos Martins, A. Liso, S. Gieowarsingh, F. Castriota

Interventional Cardio-Angiology Unit, Department of Medical and Surgical Cardiology, Villa Maria Cecilia Hospital, Via Corriera 1, 48010 Cotignola (RA), Italy

Although balloon angioplasty (BA) has been extensively reported to treat peripheral artery occlusive disease, only little information is available about its use for critical hand ischemia. We report a case

of a patient with painful ulcerated ischemic lesions of the right hand digits successfully treated by BA of the radial artery (RA) and its deep palmar branch (DPB). The 6-month angiographic follow-up showed a focal restenosis that was approached by a second BA session. Larger series are encouraged to determine the clinical outcome of percutaneous approach in this arterial district.

doi:10.1016/j.ejvs.2009.01.022

DOI of original article:10.1016/j.ejvs.2009.01.003

Available online 26 March 2009

### Behcet's Disease presenting as Bilateral Common Femoral Artery Pseudo-aneurysms

G.W.L. Tan<sup>\*</sup>, K.H. Chia

Vascular Surgery Service, Department of General Surgery, Tan Tock Seng Hospital, Singapore

We present a 25-year-old Chinese male who was diagnosed with Behcet's disease after suffering bilateral femoral artery pseudo-aneurysms. This was treated surgically by excising the pseudo-aneurysm and achieving vascular continuity by harvesting the long saphenous vein as a conduit. Arterial pseudo-aneurysms are rare vascular manifestations in Behcet's disease and management with vein is advisable as patients are frequently on long-term immunosuppression which puts them at risk of infections with prosthetic grafts.

doi:10.1016/j.ejvs.2009.02.011

DOI of original article:10.1016/j.ejvsextra.2009.02.001

Available online 31 March 2009

<sup>☆</sup> Full articles available online at [www.ejvsextra.com](http://www.ejvsextra.com)